arXiv:1304.5495v1 [math-ph] 19 Apr 2013

Harmonic and Dirac oscillators in a (2+1)-dimensional noncommutative space

F. Vega*
IFLP - CONICET and Departamento de Física,
Facultad de Ciencias Exactas de la UNLP,
C.C. 67, (1900) La Plata, Argentina
(Dated: April 22, 2013)

We study the Harmonic and Dirac Oscillator problem extended to a three-dimensional noncommutative space where the noncommutativity is induced by a shift of the dynamical variables with generators of $SL(2,\mathbb{R})$ in a unitary irreducible representation. The Hilbert space gets the structure of a direct product with the representation space as a factor, where there exist operators which realize the algebra of Lorentz transformations.

The spectrum of these models are considered in perturbation theory, both for small and large noncommutativity parameters, finding no constraints between coordinates and momenta noncommutativity parameters.

Since the representation space of the unitary irreducible representations $SL(2,\mathbb{R})$ can be realized in terms of spaces of square-integrable functions, we conclude that these models are equivalent to quantum mechanical models of particles living in a space with an additional compact dimension.

I. INTRODUCTION

In reference [1], a (2+1)-dimensional model with a kind of nonstandard noncommutativity has been considered, where both coordinates and momenta get deformed commutators. In order to preserve the Lorentz invariance, these deformations of the Heisenberg algebra were taken as proportional to generators of the Lorentz group in some irreducible representation (*irrep*) of the (noncompact) Lie group $SL(2,\mathbb{R})$.

This deformed algebra can be effectively realized by shifting the ordinary (commutative) dynamical variables by generators of the irreducible representation of $SL(2,\mathbb{R})$ considered. This means that the Hilbert space has the structure of a direct product, where one factor corresponds to the component of the state vectors in the representation space of the *irrep* considered.

As we demand that the noncommuting phase space variables to be Hermitian operators, we are constrained to consider unitary *irrep*'s of this group, which are not of finite dimension. Moreover, since the representation space of the unitary *irrep*'s of $SL(2,\mathbb{R})$ can be explicitly realized in terms of spaces of functions defined on the unit circle or analytic functions on the unit open disk (the irreducible representations of $SL(2,\mathbb{R})$ have been extensively discussed in [2]; see also [1]), the models to be considered turn out to be equivalent to quantum mechanical systems living in a space with an additional (compact) dimension.

In [1] the Landau problem for bosonic and fermionic particles living on this noncommutative space have been considered. The goal of the present article is to extend this analysis to the case of the harmonic and Dirac oscillators appropriately formulated on this space.

II. SETTING OF THE PROBLEM

According to the ideas exposed in [1], we consider the modified Heisenberg algebra of the (Hermitian) dynamical variables given by

$$\begin{aligned}
&[\hat{x}_\mu, \hat{x}_\nu] = -i\theta^2 \epsilon_{\mu\nu\rho} s^\rho \,, && [\hat{p}_\mu, \hat{p}_\nu] = -i\kappa^2 \epsilon_{\mu\nu\rho} s^\rho \,, \\
&[\hat{x}_\mu, \hat{p}_\nu] = i(\eta_{\mu\nu} - \kappa\theta \epsilon_{\mu\nu\rho} s^\rho) \,, && [\hat{x}_\mu, s_\nu] = -i\theta \epsilon_{\mu\nu\rho} s^\rho \,, \\
&[\hat{p}_\mu, s_\nu] = -i\kappa \epsilon_{\mu\nu\rho} s^\rho \,, && [s_\mu, s_\nu] = -i\epsilon_{\mu\nu\rho} s^\rho \,,
\end{aligned} \tag{1}$$

where $s_\mu \,, \mu = 0, 1, 2$ are the generators of an *irrep* of $SL(2,\mathbb{R})$ and θ and κ play the role of ultraviolet and infrared scales respectively.

---

*Electronic address: federicogaspar@gmail.com

This deformation can be realized in terms of dynamical variables satisfying the usual Heisenberg algebra, x^μ, p_μ with $\mu = 0, 1, 2$, through the *non-Abelian Bopp's shift* given by

$$\hat{x}_\mu \to x_\mu + \theta s_\mu \,, \quad \hat{p}_\mu \to p_\mu + \kappa s_\mu \,. \tag{2}$$

We follow the notation of [1], where it has been shown that the operators $M_\mu := L_\mu + s_\mu$, with $L^\mu := \frac{1}{2}\,\epsilon^{\mu\nu\lambda}\,(x_\nu p_\mu - x_\mu p_\nu)$, are a realization on the Hilbert space of the system of the generators of the Lorentz transformations in the 2+1-dimensional noncommutative space we are considering. Indeed (See ref. [1]), it can be seen that

$$\begin{aligned} [M_\mu, M_\nu] &= -\imath \epsilon_{\mu\nu\lambda} M^\lambda \quad [M_\mu, s_\nu] = -\imath \epsilon_{\mu\nu\lambda} s^\lambda \,, \\ [M_\mu, \hat{x}_\nu] &= -\imath \epsilon_{\mu\nu\lambda} \hat{x}^\lambda \,, \quad [M_\mu, \hat{p}_\nu] = -\imath \epsilon_{\mu\nu\lambda} \hat{p}^\lambda \,. \end{aligned} \tag{3}$$

In the following, our strategy to formulate a model in this noncommutative space will be, given a Hamiltonian $H(\mathbf{p}, \mathbf{x})$ in the usual (commutative) Minkowski space, to generalize it by taking $H(\hat{\mathbf{p}}, \hat{\mathbf{x}})$ and then analyze it through the replacements in Eq. (2). In this way, we construct models of Schrödinger and Dirac oscillators in this NC extension of the 2+1-dimensional Minkowski phase space and consider the implications this generalization may have.

In particular, as already mentioned, we get a space of state vectors which is the direct product of the Hilbert space for the systems in the usual commutative space with the representation space of a unitary *irrep* of $SL(2, \mathbb{R})$. Since these representation spaces can be realized as spaces of square-integrable functions (functions on the unit circle or analytic functions on the open unit disk, according to the particular *irrep* considered), these models can also be interpreted as describing particles living in spaces with an additional (compact) spatial dimension.

III. HARMONIC OSCILLATOR

A. The usual case

Let us first consider the Schrödinger Hamiltonian for the harmonic oscillator:

$$2MH := p_i^2 + M^2\omega^2 x_i^2 \,, \tag{4}$$

which commutes with the generator of rotations on the plane, L_0.

In terms of creation and annihilation operators (see [1] for the notation) the Hamiltonian H and the angular momentum L_0 can be written as

$$H := \omega(a^\dagger a + b^\dagger b + 1), \quad L_0 := b^\dagger b - a^\dagger a \,. \tag{5}$$

Their simultaneous eigenvectors are

$$|n_a, n_b\rangle := \frac{\left(a^\dagger\right)^{n_a}}{\sqrt{n_a!}} \, \frac{\left(b^\dagger\right)^{n_b}}{\sqrt{n_b!}} \, |0, 0\rangle \;, \quad n, N = 0, 1, 2, \ldots$$

where $a\,|0, 0\rangle = 0 = b\,|0, 0\rangle$, corresponding to the eigenvalues

$$H\,|n_a, n_b\rangle = \omega\,(n_a + n_b + 1)\,|n_a, n_b\rangle \;, \quad L_0\,|n_a, n_b\rangle = (n_b - n_a)\,|n_a, n_b\rangle \;. \tag{6}$$

We call $l := (n_b - n_a)$.

B. Extension to the noncommutative space

The generalization of this system to the noncommutative space defined by Eq. (1) is

$$2M\hat{H} := \hat{p}_i^2 + M^2\omega^2 \hat{x}_i^2 = (p_i + \kappa s_i)^2 + M^2\omega^2\,(x_i + \theta s_i)^2 \;, \tag{7}$$

which commutes with $M_0 = L_0 + s_0$, as can be easily seen..

In terms of creation and annihilation operators and the Hermitian generators s_μ, this Hamiltonian reads as

$$2M\hat{H} = 2MH + \sqrt{M\omega}\left(za^\dagger s_+ + \overline{z}as_- + zb^\dagger s_- + \overline{z}bs_+\right) + z\overline{z}\left({s_0}^2 - \mathbf{s}^2\right) , \tag{8}$$

where $z := \theta M\omega + \imath\kappa$, $\overline{z} := \theta M\omega - \imath\kappa$ and we have defined

$$s_\pm := s_1 \pm \imath s_2 , \quad \mathbf{s}^2 := {s_0}^2 - {s_1}^2 - {s_2}^2 . \tag{9}$$

In [1] one can find a brief review of the unitary irreducible representations (*irrep*) of $sl(2,\mathbb{R})$. The representation space is generated by the basis of simultaneous eigenvectors of s_0 and $\mathbf{s}^2$,

$$\mathbf{s}^2 \left|\lambda, m\right\rangle = \lambda \left|\lambda, m\right\rangle , \quad s_0 \left|\lambda, m\right\rangle = m \left|\lambda, m\right\rangle , \tag{10}$$

where λ and m are real numbers. We also have

$$s_\pm \left|\lambda, m\right\rangle = \sqrt{m(m\pm 1) - \lambda}\, \left|\lambda, m\pm 1\right\rangle . \tag{11}$$

The *discrete classes* of unitary *irrep*'s [1, 2] correspond to $\lambda \geq -1/4$, for which this parameter takes discrete values, $\lambda = k(k-1)$ with $k = \frac{N}{2}, N \in \mathbb{N}$, and either $m = k, k+1, k+2, \cdots$ or $m = -k, -k-1, -k-2, \cdots$ On the other hand, for the *continuous classes* of unitary *irrep*'s [1, 2], λ takes any real value less than $-1/4$ and m takes either all the integer or all the half-integer values.

The Hilbert space is then the linear span of the vectors of the form

$$\left|n_a, n_b; \lambda, m\right\rangle := \left|n_a, n_b\right\rangle \otimes \left|\lambda, m\right\rangle , \tag{12}$$

which are simultaneously eigenvectors of H, L_0, $\mathbf{s}^2$ and s_0, normalized so as to satisfy

$$\left\langle n_a, n_b; \lambda, m | n_a', n_b'; \lambda, m'\right\rangle = \delta_{n_a,n_a'}\delta_{n_b,n_b'}\delta_{m,m'} . \tag{13}$$

We also have that $\left[\hat{H}, M_0\right] = 0$, where $M_0 = L_0 + s_0$ has eigenvalues $j = l + m = n_b - n_a + m$, integer or half-integer according to the *irrep* of $SL(2,\mathbb{R})$ considered. Then, for given values of λ and j, we can give the following development for the $\hat{H}$'s eigenvectors,

$$\left|\psi_{E,j}\right\rangle = \sum_{n_b - n_a + m = j} C_{n_a,n_b,m} \left|n_a, n_b; \lambda, m\right\rangle . \tag{14}$$

From Eq. (8), one straightforwardly gets the linear recursion relation for the coefficients

$$\begin{gathered}\left\langle n_a, n_b; \lambda, m\right| 2M(\hat{H} - E) \left|\psi_{E,j,n_b}\right\rangle = \left\{2M\omega(n_a + n_b + 1) - 2M(E - \kappa m) + \bar{z}z\left(m^2 - \lambda\right)\right\} C_{n_a,n_b,m} + \\ + z\sqrt{M\omega}\sqrt{n_a+1}\sqrt{m(m+1) - \lambda}\, C_{n_a+1,n_b,m+1} + \bar{z}\sqrt{M\omega}\sqrt{n_a}\sqrt{(m-1)m - \lambda}\, C_{n_a-1,n_b,m-1} + \\ + z\sqrt{M\omega}\sqrt{n_b+1}\sqrt{m(m-1) - \lambda}\, C_{n_a,n_b+1,m+1} + \bar{z}\sqrt{M\omega}\sqrt{n_b}\sqrt{(m+1)m - \lambda}\, C_{n_a,n_b-1,m-1} = 0 ,\end{gathered} \tag{15}$$

where $m = j + n_a - n_b$.

Notice that, for $z = 0$, this recurrence gives immediately the usual harmonic oscillator levels,

$$C_{n,m}\left[\omega(n_a + n_b + 1) - E\right] = 0 \quad \Rightarrow \quad E = \omega\left(n_a + n_b + 1\right) . \tag{16}$$

It is not evident how to get an exact solution of this recurrence for $z \neq 0$. A difference with the Landau problem studied in [1] is that $\left[H, b^\dagger b\right] \neq 0$. A direct consequence of this, is that the eigenvalues problem does not reduce to a matricial one for unitary *irrep*'s in the discrete classes.

C. The spectrum in perturbation theory

1. *Small $|z|$*

In order to explain the structure of this spectrum we will use perturbation theory for small values of the noncommutativity parameters. For convenience, we take as unperturbed Hamiltonian H_0 and perturbation V the operators given by

$$
\begin{aligned}
2MH_0 &= 2M\omega\left[a^\dagger a + b^\dagger b + 1\right] + 2M\kappa s_0 + \bar{z}z\left({s_0}^2 - \mathbf{s}^2\right) \,, \\
2MV &= \sqrt{M\omega}\left(za^\dagger s_+ + \overline{z}as_- + zb^\dagger s_- + \overline{z}bs_+\right) \,.
\end{aligned} \tag{17}
$$

Since H_0 commutes with L_0 and s_0, the unperturbed eigenvectors and eigenvalues are given by

$$
\begin{aligned}
\Psi_{n_a,n_b,m} = |n_a, n_b\rangle \otimes |\lambda, m\rangle \,, \quad H_0\Psi_{n_a,n_b,m} = E^{(0)}_{n_a,n_b,m}\Psi_{n_a,n_b,m} \,, \\
E^{(0)}_{n_a,n_b,m} = \omega\left(n_a + n_b + 1\right) + \kappa m + \frac{\bar{z}z}{2M}\left(m^2 - \lambda\right) \,.
\end{aligned} \tag{18}
$$

The first order corrections to the eigenvalues in perturbation theory are simply given by

$$
E^{(1)}_{n_a,n_b,m} = \left(\Psi_{n_a,n_b,m}, V\Psi_{n_a,n_b,m}\right) = 0 \,, \tag{19}
$$

and are all vanishing.

The second order corrections are given by

$$
E^{(2)}_{n_a,n_b,m} = {\sum_{n'_a,n'_b,m'}}' \frac{\left|\left(\Psi_{n'_a,n'_b,m'}, V\Psi_{n_a,n_b,m}\right)\right|^2}{E^{(0)}_{n_a,n_b,m} - E^{(0)}_{n'_a,n'_b,m'}} \,, \tag{20}
$$

where the term with $n'_a = n_a$, $n'_b = n_b$ and $m' = m$ is excluded from the series. From (17) we get

$$
\begin{aligned}
&\left(\Psi_{n'_a,n'_b,m'}, 2MV\Psi_{n_a,n_b,m}\right) = \\
&= \sqrt{M\omega}z\sqrt{n_a+1}\sqrt{m(m+1)-\lambda}\,\delta_{n'_a,n_a+1}\delta_{n'_b,n_b}\delta_{m',m+1} + \sqrt{M\omega}\bar{z}\sqrt{n_a}\sqrt{m(m-1)-\lambda}\,\delta_{n'_a,n_a-1}\delta_{n'_b,n_b}\delta_{m',m-1} + \\
&+ \sqrt{M\omega}z\sqrt{n_b+1}\sqrt{m(m-1)-\lambda}\,\delta_{n'_a,n_a}\delta_{n'_b,n_b+1}\delta_{m',m-1} + \sqrt{M\omega}\bar{z}\sqrt{n_b}\sqrt{m(m+1)-\lambda}\,\delta_{n'_a,n_a}\delta_{n'_b,n_b-1}\delta_{m',m+1} \,,
\end{aligned} \tag{21}
$$

from which it follows that

$$
E^{(2)}_{n_a,l,m} = -\frac{|z|^2}{2M}\left(m^2 - \lambda\right) - \frac{|z|^2}{2M}ml + O\left(|z|^3\right) \,. \tag{22}
$$

Then, up to second order in $|z|$, we get for the eigenvalues

$$
E_{n_a,n_b,m} = \omega\left(n_a + n_b + 1\right) + \kappa m - \frac{|z|^2}{2M}ml + O\left(|z|^3\right) \,, \tag{23}
$$

for any unitary *irrep* of $SL(2,\mathbb{R})$. Notice that the linear in κ correction produces, for each given m, a rigid shift of the (zero order) harmonic oscillator levels. Moreover, the term quadratic in $|z|^2$ shows a coupling with the angular momentum, thus breaking the usual degeneracy in l of the isotropic harmonic oscillator spectrum. Notice also that the dominant term in the θ parameter is quadratic.

2. Large $|z|$

We will also consider the large NC parameters limit in perturbation theory. In this limit we consider the operator in Eq. 8 divided by $|z|^2$ and take as unperturbed Hamiltonian the operator

$$
\frac{\mathcal{H}_0}{|z|^2} := \frac{1}{2M}\left({s_0}^2 - \mathbf{s}^2\right) + \frac{\kappa}{|z|^2}s_0 + \frac{\omega}{|z|^2}\left[a^\dagger a + b^\dagger b + 1\right] \,, \tag{24}
$$

and as perturbation

$$
\frac{\mathcal{V}}{|z|^2} := \sqrt{\frac{\omega}{M}}\,\frac{1}{|z|^2}\left(za^\dagger s_+ + \overline{z}as_- + zb^\dagger s_- + \overline{z}bs_+\right) \,. \tag{25}
$$

The eigenvectors and eigenvalues of $\mathcal{H}_0$ are given by

$$\chi_{n_a,n_b,m} := |n_a,n_b\rangle \otimes |\lambda, m\rangle \ ,$$

$$\frac{\mathcal{E}^{(0)}_{n_a,n_b,m}}{|z|^2} = \frac{1}{2M}\left(m^2-\lambda\right) + \frac{\kappa}{|z|^2}\, m + \frac{\omega}{|z|^2}(n_a+n_b+1)\,, \tag{26}$$

while the first order correction to the eigenvalues in perturbation theory vanish,

$$(\chi_{n_a,n_b,m}, \mathcal{V}\chi_{n_a,n_b,m}) = 0\,. \tag{27}$$

Therefore, in the large $|z|$ limit, the noncommutativity parameters appear as a typical energy scale for the separation of successive series of isotropic harmonic oscillator levels. For $|z|/M \gg 1$, only the states with the minimum value of m^2 will manifest al low energies.

IV. DIRAC OSCILLATOR

The free Dirac equation in 2+1-dimensions is

$$(\imath\gamma^\mu \partial_\mu - M)\,\Psi = 0\,, \tag{28}$$

where we take

$$\gamma^0 = \sigma_3\,, \quad \gamma^1 = -\imath\sigma_2\,, \quad \gamma^2 = \imath\sigma_1\,, \tag{29}$$

which satisfy $[\gamma^\mu,\gamma^\nu] = 2g^{\mu\nu}$ with $(g^{\mu\nu}) = \mathrm{diag}\,(1,-1,-1)$. From (28) we get the Hamiltoniano $H = \alpha_i \mathbf{p}_i + M\beta$, where $\alpha_1 = -\sigma_1$, $\alpha_2 = -\sigma_2$, $\beta = \sigma_3$ and $M > 0$ the mass of the particle.

In [3] Moshinsky and Szczepaniak proposed to add a linear in the coordinates term, interpreting the resulting system as a *Dirac oscillator* since, in the nonrelativistic limit, it reduces to a harmonic oscillator with a spin-orbit interaction. In this way, the Hamiltonian operator becomes

$$H = \alpha_i\,(p_i - \imath\omega\beta x_i) + M\beta\,, \tag{30}$$

for some constant $\omega > 0$.

In terms of creation and annihilation operators (see [1] for the notation, identifying $\omega \leftrightarrow eB/2$) the Hamiltonian H and the angular momentum L_0 read now as

$$H = \begin{pmatrix} M & \sqrt{4\omega}a^\dagger \\ \sqrt{4\omega}a & -M \end{pmatrix}\,, \quad L_0 = b^\dagger b - a^\dagger a\,. \tag{31}$$

A simple calculation shows H commutes with $J_0 := L_0 + \frac{\sigma_3}{2}$. The eigenvectors of H with angular momentum $j_0 = n_b - n_a + \frac{1}{2}$ are [3] (see also [4])

$$\psi_{\pm,n_a,n_b} = \begin{pmatrix} \sqrt{4\omega n_a}\,|n_a,n_b\rangle \\ \\ \left[-M \pm \sqrt{M^2+4\omega n_a}\right]|n_a-1,n_b\rangle \end{pmatrix}\,, \tag{32}$$

for $n_a \geq 1$, corresponding to the eigenvalues $E_{\pm,n_a,n_b} = \pm\sqrt{M^2+4\omega n_a}$, degenerate in the index n_b. There is another solution for $n_a = 0$, given by

$$\psi_{0,n_b} = \begin{pmatrix} |0,n_b\rangle \\ 0 \end{pmatrix}\,, \tag{33}$$

with $j_0 = n_b + 1/2$ and $E_{0,n_b} = M$, also degenerate in n_b.

A. Extension to the noncommutative space

It is easy to see that a similar analysis as that for the Dirac case in [1] can be done for the present problem. Indeed, if we adopt as Hamiltonian the Hermitian operator

$$\hat{H} = \alpha_i \left(\hat{p}_i - \imath\omega\beta\hat{x}_i \right) + \kappa s_0 + M\beta \tag{34}$$

and use the identity $\alpha_i\beta = -\imath\epsilon_{ij}\alpha_j$, we get

$$\hat{H} = H \otimes \mathbf{1} + \kappa \left(\alpha_i \otimes s_i + \mathbf{1}_2 \otimes s_0 \right) - \theta\omega\, \epsilon_{ij}\alpha_j \otimes s_i \,, \tag{35}$$

where H is given in Eqs. (30) and (31). This coincides exactly with the Hamiltonian in Eq. (60) of [1].

Therefore, the identification $\omega \equiv eB/2$ shows that the results obtained in [1] for the Landau problem for Dirac particles in this noncommutative space apply also to the extension of the Dirac oscillator presented here. We will not reproduce these results here and refer the reader to Section IV of [1].

V. CONCLUSIONS

In this article we have considered models of Schrödinger and Dirac oscillators in a space-time with a kind of noncommutativity, both in coordinates and momenta, induced by deforming the canonical commutators by terms proportional to the generators in a unitary irreducible representation of the Lorentz group in the 2+1-dimensional Minkowski space, isomorphic to $SL(2,\mathbb{R})/\mathbb{Z}_2$. Since this is a noncompact Lie group, its unitary *irrep*'s are not of finite dimension. This noncommutative phase space has been considered in [1].

This deformed Heisenberg algebra can be realized by means of a shift in the canonical coordinates and momenta with terms proportional to the generators of $SL(2,\mathbb{R})/\mathbb{Z}_2$ in a unitary *irrep*. In particular, the shift in momenta can also be interpreted as the introduction of a non-Abelian magnetic field.

Consequently, the number of dynamical variables is enlarged and the Hilbert space gets the structure of a direct product, one factor for the state vectors of the ordinary system in the normal space and the other for the component of the state vectors in the representation space of this *irrep* of $SL(2,\mathbb{R})$.

It was shown in [1] that total generators of the Lorentz transformations can be constructed which correctly transform all the operators, thus realizing the Lie algebra $sl(2,\mathbb{R})$ on the Hilbert space of these quantum-mechanical system.

In this framework, we have considered modified Hamiltonians obtained through this non-Abelian Bopp's shift of the dynamical variables from the Hamiltonians of the Harmonic and Dirac Oscillators. We have analyzed these models for both discrete and continuous classes of *irrep*'s of $SL(2,\mathbb{R})$, getting linear infinite recursion relations for the coefficients in the development of the Hamiltonian eigenvectors in terms of conveniently chosen bases of the Hilbert space. The spectrum of these models have been studied also in perturbation theory, both for small and large noncommutativity parameters $z = \theta M\omega + \imath\kappa$.

For small z, we have shown that the spectrum of the extension of the harmonic oscillator is the one of the isotropic two-dimensional oscillator rigidly shifted by κm, a term proportional to the eigenvalue of s_0. Moreover, second order corrections in $|z|$ break the degeneracy in the angular momentum. In particular, the dominant correction in the θ parameter is quadratic. In the large $|z|$ limit, the noncommutativity parameters appear as a typical energy scale for the separation of successive series of isotropic harmonic oscillator levels.

On the other hand, we have shown that the extension of the Dirac oscillator to this noncommutative space is completely equivalent to the extension of the Landau problem for Dirac particles studied in [1].

Let us mention that, contrary to the case of conventional NC Quantum Mechanics, we find no constraint between the parameters referring to no-commutativity in coordinates and momenta. Rather, both κ and θ play a similar role..

As discussed in [1], these models do not correspond to just a smooth deformation of the commutative ones. Rather, the Hilbert space takes the structure a of the direct product of the usual one times the representation space of an *irrep* of $SL(2,\mathbb{R})$. Since these *irrep*'s can be explicitly realized in terms of spaces of square-integrable functions (functions defined on the unit circle for the continuous classes of *irrep*'s and analytic functions on the unit open disk for the discrete classes of *irrep*'s - see [1, 2]) the examples studied in this article can also be considered as equivalent to models of quantum mechanical particles living in a space with an additional compact dimension, with $|z|$ playing the role of the inverse of a typical length.

Acknowledgements: F.V. acknowlege finantial support from CONICET, Argentina. This work was partially supported by Universidad Nacional de La Plata (Proy. 11/X615), Argentina.

[1] H. Falomir, F. Vega, J. Gamboa, F. Mendez, M. Loewe.*Phys. Rev. D* **86**, 105035 (2012)
[2] V. Bargmann, *Ann. Math.* **48**, 568 (1947).
[3] M. Moshinsky and A. Szczepaniak, J. Phys. A: Math. Gen. 22, L817 (1989).
[4] Nagalakshmi A. Rao and B.A. Kagali, arXiv:0704.3362 [quant-ph] (2007).